

Halfconvex functions and equality of Daróczy–Páles conjugate means

JANUSZ MATKOWSKI

Abstract. Using some properties of halfconvex and halfconcave functions, we solve the equality problem for conjugate means introduced by Daróczy and Páles.

Mathematics Subject Classification (2010). Primary 26A51, 26D15, 26E60.

Keywords. Mean, conjugate mean, (s, p) -halfconvex function, (s, p) -halfconvex function.

1. Introduction

For a mean $K : I^2 \rightarrow I$, a continuous strictly monotonic function φ defined in an interval $I \subset \mathbb{R}$, and the numbers $p, q \in [0, 1]$, Daróczy and Páles [3] defined a K -conjugate mean $K_\varphi^{(p,q)}$ by the formula

$$K_\varphi^{(p,q)}(x, y) := \varphi^{-1}(p\varphi(x) + q\varphi(y) + (1 - p - q)\varphi(K(x, y))), \quad x, y \in I.$$

In the case $p = q = 1$ such a mean was considered in [6].

In [2] the authors solve the equality problem of a conjugate mean and a weighted quasi-arithmetic mean. It is interesting that this equality problem contains the invariance equation, considered among others in [4, 6], as a special case.

In the present paper we consider the question of equality of conjugate means, that is, we deal with the equation $L_\psi^{(r,s)} = K_\varphi^{(p,q)}$. This problem easily reduces to the equality $L_{id}^{(r,s)} = K_\gamma^{(p,q)}$ where $\gamma := \varphi \circ \psi^{-1}$ (Sect. 3). In Sect. 4 we show that this leads to inequalities of the form

$$\gamma((1 - s)x + sy) \leq p\gamma(x) + (1 - p)\gamma(y), \quad x, y \in I, \quad x \leq y;$$

and

$$\gamma((1 - s)x + sy) \geq p\gamma(x) + (1 - p)\gamma(y), \quad x, y \in I, \quad x \leq y$$

(Lemmas 4.1–4.4). These weaker forms of (s, p) -convexity ((s, p) -concavity) of a function appeared for the first time in a recent paper of Daróczy [1], where the function satisfying the first of these inequalities is said to be (s, p) -halfconvex. Some auxiliary results on the properties of (s, p) -halfconvex and (s, p) -halfconcave functions presented in Sect. 5 allow us to prove the main result which reads as follows: *if $K, L : I^2 \rightarrow I$ are means in the interval I , $p, q, r, s \in [0, 1]$ are fixed numbers such that $p + q \neq 1 \neq r + s$, and $\varphi, \psi : I \rightarrow \mathbb{R}$ are continuous strictly monotonic functions, then $L_\psi^{(r,s)} = K_\varphi^{(p,q)}$ if, and only if, $\psi \sim \varphi$ and, for all $x, y \in I$,*

$$L(x, y) = \varphi^{-1} \left(\frac{(p-r)\varphi(x) + (q-s)\varphi(y) + (1-p-q)\varphi(K(x, y))}{1-r-s} \right).$$

2. Preliminaries

For an interval $I \subset \mathbb{R}$, by $\mathcal{CM}(I)$ denote the family of all continuous and strictly monotonic functions $\varphi : I \rightarrow \mathbb{R}$, and by $\mathcal{CM}_+(I)$ the family of all continuous and strictly increasing functions $\varphi : I \rightarrow \mathbb{R}$. The functions $\varphi, \psi \in \mathcal{CM}(I)$ are called equivalent, in writing $\psi \sim \varphi$, if $\psi = a\varphi + b$ for some $a, b \in \mathbb{R}, a \neq 0$.

Recall that a function $M : I^2 \rightarrow I$ is called a mean if

$$\min(x, y) \leq M(x, y) \leq \max(x, y), \quad x, y \in I.$$

A mean M is called strict if for all $x, y \in I, x \neq y$, these inequalities are sharp; symmetric if $M(x, y) = M(y, x)$.

Let $I \subset \mathbb{R}$ be an interval, $K : I^2 \rightarrow I$ a mean in $I, p, q \in [0, 1]$ fixed numbers and $\varphi \in \mathcal{CM}(I)$. Then (cf. [3]) the function $K_\varphi^{(p,q)} : I^2 \rightarrow \mathbb{R}$ defined by

$$K_\varphi^{(p,q)}(x, y) := \varphi^{-1}(p\varphi(x) + q\varphi(y) + (1-p-q)\varphi(K(x, y))), \quad x, y \in I,$$

is a mean and it is called the K -conjugate mean of the quasi-generator φ , and the weights p, q . Note that

$$K_\varphi^{(0,0)} = K \quad \text{and} \quad K_\varphi^{(p,1-p)} = A_p^{[\varphi]},$$

where

$$A_p^{[\varphi]}(x, y) := \varphi^{-1}(p\varphi(x) + (1-p)\varphi(y)), \quad x, y \in I,$$

is the quasi-arithmetic mean of the generator φ and the weight p (cf. [2]).

Thus the family of means $\{K_\varphi^{(p,q)} : p, q \in [0, 1]\}$ contains the mean K and all the weighted quasi-arithmetic means $A_p^{[\varphi]}$. Moreover, from the equality

$$K_\varphi^{(1,1)}(x, y) = \varphi^{-1}(\varphi(x) + \varphi(y) - \varphi(K(x, y))), \quad x, y \in I,$$

we get

$$\varphi(K(x, y)) + \varphi(K_\varphi^{(1,1)}(x, y)) = \varphi(x) + \varphi(y), \quad x, y \in I,$$

whence, setting $A^{[\varphi]} := A_{1/2}^{[\varphi]}$, we obtain

$$A^{[\varphi]} \circ (K, K_\varphi^{(1,1)}) = A^{[\varphi]},$$

that is, the quasi-arithmetic mean $A^{[\varphi]}$ is invariant with respect to the mean-type mapping $(K, K_\varphi^{(1,1)}) : I^2 \rightarrow I^2$ (cf. [6], Remark 2). This fact is important in applications.

Remark 2.1. If $\varphi, \psi \in \mathcal{CM}(I)$ and $\varphi \sim \psi$, then $K_\psi^{(p,q)} = K_\varphi^{(p,q)}$. Therefore we can always assume that the generator φ of the mean $K_\varphi^{(p,q)}$ is strictly increasing, i.e. that $\varphi \in \mathcal{CM}_+(I)$.

3. Problem and remarks

If $L : I^2 \rightarrow I$ is another mean in I , the numbers $r, s \in [0, 1]$ are fixed, and $\psi \in \mathcal{CM}(I)$, then we can ask when

$$L_\psi^{(r,s)} = K_\varphi^{(p,q)},$$

that is, when two conjugate means of different generators and weights can be equal?

The following is easy to verify.

Remark 3.1. Assume that $I \subset \mathbb{R}$ is an interval, $K, L : I^2 \rightarrow I$ are means, $p, q, r, s \in [0, 1]$ are fixed numbers, and $\varphi, \psi \in \mathcal{CM}(I)$. Then

$$L_\psi^{(r,s)} = K_\varphi^{(p,q)}$$

if, and only if,

$$\mathcal{L}_{id}^{(r,s)} = \mathcal{K}_\gamma^{(p,q)},$$

where $\mathcal{L}, \mathcal{K} : \psi(I) \times \psi(I) \rightarrow \psi(I)$ are means defined by

$$\mathcal{K}(u, v) := \psi(K(\psi^{-1}(u), \psi^{-1}(v))), \quad \mathcal{L}(u, v) := \psi(L(\psi^{-1}(u), \psi^{-1}(v)))$$

for all $u, v \in \psi(I)$, and

$$\gamma := \varphi \circ \psi^{-1};$$

that is

$$\mathcal{K}_\gamma^{(p,q)}(u, v) = \gamma^{-1}(p\gamma(u) + q\gamma(v) + (1 - p - q)\gamma(\mathcal{K}(u, v))), \quad u, v \in \psi(I),$$

and

$$\mathcal{L}_{id}^{(r,s)} = ru + sv + (1 - r - s)\mathcal{L}(u, v), \quad u, v \in \psi(I).$$

This remark allows us to reduce the problem of equality of conjugate means to the functional equation $\mathcal{L}_{id}^{(r,s)} = \mathcal{K}_\gamma^{(p,q)}$, where id denotes the identity function and γ is continuous and strictly monotonic.

Remark 3.2. Let $I \subset \mathbb{R}$ be an interval and $p, q, r, s \in [0, 1]$. Suppose that $\gamma \in \mathcal{CM}(I)$ and the functions $K, L : I^2 \rightarrow I$ are means in I .

- (i) If $p + q = 1$ and $r + s = 1$, then $L_{id}^{(r,s)} = K_\gamma^{(p,q)}$ iff $\gamma \sim id$ and $r = p, s = q$.
 Moreover, in this case both means K and L can be arbitrary.
- (ii) If $p + q = 1$ and $r + s \neq 1$, then $L_{id}^{(r,s)} = K_\gamma^{(p,q)}$ iff

$$L(x, y) = \frac{\gamma^{-1}(p\gamma(x) + (1-p)\gamma(y)) - rx - sy}{1 - r - s}, \quad x, y \in I.$$

Moreover, in this case the mean K can be arbitrary.

- (iii) If $p + q \neq 1$ and $r + s = 1$, then $L_{id}^{(r,s)} = K_\gamma^{(p,q)}$ iff

$$K(x, y) = \gamma^{-1} \left(\frac{\gamma(rx + (1-r)y) - p\gamma(x) - q\gamma(y)}{1 - p - q} \right), \quad x, y \in I.$$

Moreover, in this case the mean L can be arbitrary.

Proof. (i) Since $q = 1 - p, s = 1 - r$, by the definition of conjugate means, the equality $L_{id}^{(r,s)} = K_\gamma^{(p,q)}$ reduces to

$$rx + (1-r)y = \gamma^{-1}(p\gamma(u) + (1-q)\gamma(v)), \quad x, y \in I,$$

which holds true iff $\gamma \sim id$ and $r = p, s = q$ (cf. [5]).

The cases (ii) and (iii) follow directly from the definitions of the means $L_{id}^{(r,s)}$ and $K_\gamma^{(p,q)}$. \square

4. Equality of conjugate means implies halfconvexity and halfconcavity of quasi-generator

We begin this section with the following

Lemma 4.1. *Let $I \subset \mathbb{R}$ be an interval and $p, q, r, s \in [0, 1]$ be such that $p + q < 1$ and $r + s < 1$. Suppose that $\gamma \in \mathcal{CM}_+(I)$, the functions $K, L : I^2 \rightarrow I$ are means in I , and*

$$L_{id}^{(r,s)} = K_\gamma^{(p,q)}. \tag{1}$$

Under these assumptions, for all $x, y \in I$,

- (i) *if $x \leq y$ then*

$$\gamma((1-s)x + sy) \leq p\gamma(x) + (1-p)\gamma(y),$$

$$\gamma(rx + (1-r)y) \geq (1-q)\gamma(x) + q\gamma(y);$$

- (ii) *if $x > y$ then*

$$\gamma(rx + (1-r)y) \leq (1-q)\gamma(x) + q\gamma(y),$$

$$\gamma((1-s)x + sy) \geq p\gamma(x) + (1-p)\gamma(y).$$

Proof. If (1) holds, that is, if

$$rx + sy + (1 - r - s)L(x, y) = \gamma^{-1}(p\gamma(x) + q\gamma(y) + (1 - p - q)\gamma(K(x, y)))$$

for all $x, y \in I$, then

$$L(x, y) = \frac{\gamma^{-1}(p\gamma(x) + q\gamma(y) + (1 - p - q)\gamma(K(x, y))) - rx - sy}{1 - r - s}, \quad x, y \in I,$$

whence, for all $x, y \in I$,

$$\begin{aligned} \min(x, y) &\leq \frac{\gamma^{-1}(p\gamma(x) + q\gamma(y) + (1 - p - q)\gamma(K(x, y))) - rx - sy}{1 - r - s} \\ &\leq \max(x, y). \end{aligned} \quad (2)$$

Since $r + s < 1$ and $p + q < 1$, we get

$$\begin{aligned} &\gamma^{-1} \left(\frac{\gamma((1 - r - s)\min(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1 - p - q} \right) \\ &\leq K(x, y) \leq \gamma^{-1} \left(\frac{\gamma((1 - r - s)\max(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1 - p - q} \right) \end{aligned}$$

for all $x, y \in I$. Since K is a mean, it follows that

$$\gamma^{-1} \left(\frac{\gamma((1 - r - s)\min(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1 - p - q} \right) \leq \max(x, y)$$

for all $x, y \in I$, and

$$\gamma^{-1} \left(\frac{\gamma((1 - r - s)\max(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1 - p - q} \right) \geq \min(x, y),$$

for all $x, y \in I$, whence, as γ is strictly increasing,

$$\gamma((1 - r - s)\min(x, y) + rx + sy) \leq (1 - p - q)\gamma(\max(x, y)) + p\gamma(x) + q\gamma(y)$$

for all $x, y \in I$, and

$$\gamma((1 - r - s)\max(x, y) + rx + sy) \geq (1 - p - q)\gamma(\min(x, y)) + p\gamma(x) + q\gamma(y)$$

for all $x, y \in I$. Now, if $x < y$, from these inequalities, we get

$$\begin{aligned} \gamma((1 - s)x + sy) &\leq p\gamma(x) + (1 - p)\gamma(y), \\ \gamma(rx + (1 - r)y) &\geq (1 - q)\gamma(x) + q\gamma(y); \end{aligned}$$

and, if $x > y$, we get

$$\begin{aligned} \gamma(rx + (1 - r)y) &\leq (1 - q)\gamma(x) + q\gamma(y), \\ \gamma((1 - s)x + sy) &\geq p\gamma(x) + (1 - p)\gamma(y). \end{aligned}$$

□

In the sequel of this section we consider three remaining cases.

Lemma 4.2. *Let $I \subset \mathbb{R}$ be an interval and $p, q, r, s \in [0, 1]$ be such that $p+q > 1$ and $r+s > 1$. Suppose that $\gamma \in \mathcal{CM}(I)$, the functions $K, L : I^2 \rightarrow I$ are means in I , and*

$$L_{id}^{(r,s)} = K_{\gamma}^{(p,q)}.$$

Under these assumptions, for all $x, y \in I$,

(i) *if $x \leq y$ then*

$$\begin{aligned}\gamma((1-s)x + sy) &\geq p\gamma(x) + (1-p)\gamma(y), \\ \gamma(rx + (1-r)y) &\leq (1-q)\gamma(x) + q\gamma(y);\end{aligned}$$

(ii) *if $x > y$ then*

$$\begin{aligned}\gamma(rx + (1-r)y) &\geq (1-q)\gamma(x) + q\gamma(y), \\ \gamma((1-s)x + sy) &\leq p\gamma(x) + (1-p)\gamma(y).\end{aligned}$$

Proof. Similarly as in the proof of Lemma 4.1 we can assume that γ is an increasing function. Calculating $L(x, y)$ from the equality $L_{id}^{(r,s)} = K_{\gamma}^{(p,q)}$ and taking into account that $\min(x, y) \leq L(x, y) \leq \max(x, y)$, we obtain inequalities (2) for all $x, y \in I$. Since $r+s > 1$ and $p+q > 1$, we get

$$\begin{aligned}\gamma^{-1} \left(\frac{\gamma((1-r-s)\min(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right) \\ \leq K(x, y) \leq \gamma^{-1} \left(\frac{\gamma((1-r-s)\max(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right)\end{aligned}$$

for all $x, y \in I$. Since K is a mean, it follows that

$$\gamma^{-1} \left(\frac{\gamma((1-r-s)\min(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right) \leq \max(x, y)$$

for all $x, y \in I$, and

$$\gamma^{-1} \left(\frac{\gamma((1-r-s)\max(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right) \geq \min(x, y),$$

for all $x, y \in I$. Hence, as $1-p-q < 0$, we get

$$\gamma((1-r-s)\min(x, y) + rx + sy) \geq (1-p-q)\gamma(\max(x, y)) + p\gamma(x) + q\gamma(y),$$

$x, y \in I$, and

$$\gamma((1-r-s)\max(x, y) + rx + sy) \leq (1-p-q)\gamma(\min(x, y)) + p\gamma(x) + q\gamma(y),$$

$x, y \in I$. Now, if $x < y$, from these inequalities, we get

$$\begin{aligned}\gamma((1-s)x + sy) &\geq p\gamma(x) + (1-p)\gamma(y), \\ \gamma(rx + (1-r)y) &\leq (1-q)\gamma(x) + q\gamma(y);\end{aligned}$$

and, if $x > y$, we get

$$\begin{aligned}\gamma(rx + (1-r)y) &\geq (1-q)\gamma(x) + q\gamma(y), \\ \gamma((1-s)x + sy) &\leq p\gamma(x) + (1-p)\gamma(y).\end{aligned}$$

□

Lemma 4.3. *Let $I \subset \mathbb{R}$ be an interval and $p, q, r, s \in [0, 1]$ be such that $p+q < 1$ and $r+s > 1$. Suppose that $\gamma \in \mathcal{CM}_+(I)$, the functions $K, L : I^2 \rightarrow I$ are means in I , and*

$$L_{id}^{(r,s)} = K_{\gamma}^{(p,q)}.$$

Under these assumptions, for all $x, y \in I$,

(i) *if $x \leq y$ then*

$$\begin{aligned}\gamma(rx + (1-r)y) &\leq p\gamma(x) + (1-p)\gamma(y), \\ \gamma((1-s)x + sy) &\geq (1-q)\gamma(x) + q\gamma(y);\end{aligned}$$

(ii) *if $x > y$ then*

$$\begin{aligned}\gamma((1-s)x + sy) &\leq (1-q)\gamma(x) + q\gamma(y), \\ \gamma(rx + (1-r)y) &\geq p\gamma(x) + (1-p)\gamma(y).\end{aligned}$$

Proof. Similarly as in the proofs of the previous lemmas, assuming that γ is increasing, calculating the mean L from equality (1), and taking into account that $\min(x, y) \leq L(x, y) \leq \max(x, y)$, we obtain inequalities (2) for all $x, y \in I$. Since $r+s > 1$ and $p+q < 1$, we get

$$\begin{aligned}\gamma^{-1} \left(\frac{\gamma((1-r-s)\max(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right) \\ \leq K(x, y) \leq \gamma^{-1} \left(\frac{\gamma((1-r-s)\min(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right)\end{aligned}$$

for all $x, y \in I$.

Since K is a mean, it follows that

$$\gamma^{-1} \left(\frac{\gamma((1-r-s)\max(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right) \leq \max(x, y)$$

and

$$\gamma^{-1} \left(\frac{\gamma((1-r-s)\min(x, y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right) \geq \min(x, y),$$

for all $x, y \in I$. Hence, as γ is increasing and $1-p-q > 0$, we get

$$\gamma((1-r-s)\max(x, y) + rx + sy) \leq (1-p-q)\gamma(\max(x, y)) + p\gamma(x) + q\gamma(y),$$

$x, y \in I$, and

$$\gamma((1-r-s)\min(x, y) + rx + sy) \geq (1-p-q)\gamma(\min(x, y)) + p\gamma(x) + q\gamma(y),$$

$x, y \in I$. Now, if $x < y$ from these inequalities we get

$$\begin{aligned}\gamma(rx + (1-r)y) &\leq p\gamma(x) + (1-p)\gamma(y), \\ \gamma((1-s)x + sy) &\geq (1-q)\gamma(x) + q\gamma(y);\end{aligned}$$

and, if $x > y$ we get

$$\begin{aligned}\gamma((1-s)x + sy) &\leq (1-q)\gamma(x) + q\gamma(y), \\ \gamma(rx + (1-r)y) &\geq p\gamma(x) + (1-p)\gamma(y).\end{aligned}$$

□

Lemma 4.4. *Let $I \subset \mathbb{R}$ be an interval and $p, q, r, s \in [0, 1]$ be such that $p+q > 1$ and $r+s < 1$. Suppose that $\gamma \in \mathcal{CM}_+(I)$, the functions $K, L : I^2 \rightarrow I$ are means in I , and*

$$L_{id}^{(r,s)} = K_{\gamma}^{(p,q)}.$$

Under these assumptions, for all $x, y \in I$,

(i) *if $x \leq y$ then*

$$\begin{aligned}\gamma((1-s)x + sy) &\leq p\gamma(x) + (1-p)\gamma(y) \\ \gamma(rx + (1-r)y) &\geq (1-q)\gamma(x) + q\gamma(y);\end{aligned}$$

(ii) *if $x > y$ then*

$$\begin{aligned}\gamma(rx + (1-r)y) &\geq (1-q)\gamma(x) + q\gamma(y), \\ \gamma((1-s)x + sy) &\leq p\gamma(x) + (1-p)\gamma(y).\end{aligned}$$

Proof. Assume that γ is increasing. Calculating the mean L from the assumed equality, we obtain inequalities (2) for all $x, y \in I$. Since $r+s < 1$ and $p+q > 1$, we get

$$\begin{aligned}\gamma^{-1} \left(\frac{\gamma((1-r-s)\min(x,y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right) \\ \leq K(x,y) \leq \gamma^{-1} \left(\frac{\gamma((1-r-s)\max(x,y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right)\end{aligned}$$

for all $x, y \in I$. Since K is a mean, it follows that

$$\gamma^{-1} \left(\frac{\gamma((1-r-s)\min(x,y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right) \leq \max(x,y)$$

and

$$\gamma^{-1} \left(\frac{\gamma((1-r-s)\max(x,y) + rx + sy) - p\gamma(x) - q\gamma(y)}{1-p-q} \right) \geq \min(x,y),$$

for all $x, y \in I$. Hence, as γ is increasing and $1-p-q < 0$, we get

$$\gamma((1-r-s)\max(x,y) + rx + sy) \geq (1-p-q)\gamma(\max(x,y)) + p\gamma(x) + q\gamma(y),$$

$x, y \in I$, and

$$\gamma((1-r-s)\min(x, y) + rx + sy) \leq (1-p-q)\gamma(\min(x, y)) + p\gamma(x) + q\gamma(y),$$

$x, y \in I$. Now, if $x < y$ from these inequalities we get

$$\gamma(rx + (1-r)y) \geq p\gamma(x) + (1-p)\gamma(y),$$

$$\gamma((1-s)x + sy) \leq (1-q)\gamma(x) + q\gamma(y);$$

and, if $x > y$ we get

$$\gamma((1-s)x + sy) \geq (1-q)\gamma(x) + q\gamma(y),$$

$$\gamma(rx + (1-r)y) \leq p\gamma(x) + (1-p)\gamma(y).$$

□

5. Auxiliary results on half-convex and half-concave functions

Lemma 5.1. *Let $I \subset \mathbb{R}$ be an interval and let $a, b \in (0, 1)$.*

(i) *If $f : I \rightarrow \mathbb{R}$ satisfies the inequality*

$$f(ax + (1-a)y) \leq bf(x) + (1-b)f(y), \quad x, y \in I, \quad x \leq y, \quad (3)$$

then

$$f\left(\sum_{j=0}^n \binom{n}{j} a^{n-j}(1-a)^j x_j\right) \leq \sum_{j=0}^n \binom{n}{j} b^{n-j}(1-b)^j f(x_j) \quad (4)$$

for all $n \in \mathbb{N}$ and $x_0, x_1, \dots, x_n \in I$ $x_0 \leq x_1 \leq \dots \leq x_n$.

Moreover, if (3) is satisfied for all $x, y \in I$, then (4) holds for all $n \in \mathbb{N}$ and $x_0, x_1, \dots, x_n \in I$.

(ii) *If $f : I \rightarrow \mathbb{R}$ satisfies the inequality*

$$f(ax + (1-a)y) \leq bf(x) + (1-b)f(y), \quad x, y \in I, \quad x \geq y, \quad (5)$$

then

$$f\left(\sum_{j=0}^n \binom{n}{j} a^{n-j}(1-a)^j x_j\right) \leq \sum_{j=0}^n \binom{n}{j} b^{n-j}(1-b)^j f(x_j) \quad (6)$$

for all $n \in \mathbb{N}$ and $x_0, x_1, \dots, x_n \in I$ $x_0 \geq x_1 \geq \dots \geq x_n$.

Moreover, if (5) is satisfied for all $x, y \in I$, then (6) holds for all $n \in \mathbb{N}$ and $x_0, x_1, \dots, x_n \in I$.

Proof. It is enough to prove (i). By (3) inequality (4) holds true for $n = 1$. Assume that (4) holds true for some $n \in \mathbb{N}$ and take arbitrary $x_0, x_1, \dots, x_{n+1} \in I, x_0 \leq x_1 \leq \dots \leq x_{n+1}$. Applying first inequality (4) with x_j replaced by $ax_j + (1-a)x_{j+1}$ and then inequality (3), we obtain

$$\begin{aligned}
& f \left(\sum_{j=0}^{n+1} \binom{n+1}{j} a^{n+1-j} (1-a)^j x_j \right) \\
&= f \left(\sum_{j=0}^n \binom{n}{j} a^{n-j} (1-a)^j (ax_j + (1-a)x_{j+1}) \right) \\
&\leq \sum_{j=0}^n \binom{n}{j} b^{n-j} (1-b)^j f(ax_j + (1-a)x_{j+1}) \\
&\leq \sum_{j=0}^n \binom{n}{j} b^{n-j} (1-b)^j [bf(x_j) + (1-b)f(x_{j+1})] \\
&= \sum_{j=0}^{n+1} \binom{n+1}{j} b^{n+1-j} (1-b)^j f(x_j),
\end{aligned}$$

and induction completes the proof. \square

Remark 5.2. Obviously, if inequality (3) (or (5)) is reversed, the counterpart of this lemma remains true.

Lemma 5.3. *Let $a \in (0, 1)$ be fixed. Then the set*

$$A := \left\{ \sum_{j=0}^k \binom{n}{j} a^{n-j} (1-a)^j : n \in \mathbb{N}; k \in \{0, 1, \dots, n\} \right\}$$

is dense in $[0, 1]$.

Proof. Note that $A = \bigcup_{n \in \mathbb{N}} A_n$ where, for each $n \in \mathbb{N}$,

$$A_n = \left\{ \sum_{j=0}^k \binom{n}{j} a^{n-j} (1-a)^j : k \in \{0, 1, \dots, n\} \right\}$$

forms a finite increasing sequence with the first value being equal to a^n (tending to 0 as $n \rightarrow \infty$), the last value 1, and the distance between two successive values:

$$\sum_{j=0}^k \binom{n}{j} a^{n-j} (1-a)^j - \sum_{j=0}^{k-1} \binom{n}{j} a^{n-j} (1-a)^j = \binom{n}{k} a^{n-k} (1-a)^k$$

tending to 0 as $n \rightarrow \infty$; it easily follows from the fact that, for any $k \in \mathbb{N}$,

$$\lim_{n \rightarrow \infty} \frac{\binom{n+1}{k} a^{n+1-k} (1-a)^k}{\binom{n}{k} a^{n-k} (1-a)^k} = a < 1.$$

\square

Lemma 5.4. *Let $I \subset \mathbb{R}$ be an interval and $p, s \in (0, 1)$ be fixed. Suppose that $\gamma : I \rightarrow \mathbb{R}$ is a continuous function.*

(i) *If for all $x, y \in I, x \leq y$,*

$$\gamma((1-s)x + sy) \leq p\gamma(x) + (1-p)\gamma(y), \quad (7)$$

then there exists a function $\lambda : [0, 1] \rightarrow [0, 1]$, such that for all $x, y \in I, x \geq y$,

$$\gamma(tx + (1-t)y) \leq \lambda(t)\gamma(x) + (1-\lambda(t))\gamma(y). \quad (8)$$

(ii) *If for all $x, y \in I, x \geq y$,*

$$\gamma((1-s)x + sy) \geq p\gamma(x) + (1-p)\gamma(y), \quad (9)$$

then there exists a function $\lambda : [0, 1] \rightarrow [0, 1]$, such that for all $x, y \in I, x \geq y$,

$$\gamma(tx + (1-t)y) \geq \lambda(t)\gamma(x) + (1-\lambda(t))\gamma(y). \quad (10)$$

(iii) *If inequality (7) is satisfied for all $x, y \in I, x \leq y$, and inequality (9) is satisfied for all $x, y \in I, x \geq y$, then there exists a function $\lambda : [0, 1] \rightarrow [0, 1], \lambda([0, 1]) = [0, 1]$, such that inequality (8) is satisfied for all $x, y \in I, x \leq y$; and inequality (10) is satisfied for all $x, y \in I, x \geq y$.*

Proof. (i) Setting $a := 1-s$ and $b := p$ we see that the function $f := \gamma$ satisfies inequality (3). By Lemma 5.1(i) we get

$$\gamma\left(\sum_{j=0}^n \binom{n}{j} a^{n-j}(1-a)^j x_j\right) \leq \sum_{j=0}^n \binom{n}{j} b^{n-j}(1-b)^j \gamma(x_j)$$

for all $n \in \mathbb{N}$ and $x_0, x_1, \dots, x_n \in I, x_0 \leq x_1 \leq \dots \leq x_n$.

Take arbitrary $x, y \in I, x < y$, and $k, n \in \mathbb{N}, k \leq n$. Setting here

$$x_0 = x_1 = \dots = x_k := x; \quad x_{k+1} = x_{k+2} = \dots = x_n := y$$

we obtain

$$\gamma(a_{n,k}x + (1-a_{n,k})y) \leq b_{n,k}\gamma(x) + (1-b_{n,k})\gamma(y),$$

where

$$a_{n,k} := \left(\sum_{j=0}^k \binom{n}{j} a^{n-j}(1-a)^j\right), \quad b_{n,k} := \left(\sum_{j=0}^k \binom{n}{j} b^{n-j}(1-b)^j\right).$$

By Lemma 5.3, the sets $A := \{a_{n,k} : n \in \mathbb{N}, k = 0, \dots, n\}$ and $B := \{b_{n,k} : n \in \mathbb{N}, k = 0, \dots, n\}$ are dense in $(0, 1)$. Hence, for arbitrary $t \in (0, 1)$, there is a sequence $t_j \in A, t_j = a_{n_j, k_j}, j \in \mathbb{N}$, such that $\lim_{j \rightarrow \infty} t_j = t$ and

$$\gamma(t_j x + (1-t_j)y) \leq b_{n_j, k_j}\gamma(x) + (1-b_{n_j, k_j})\gamma(y).$$

Setting a convergent subsequence (b_{j_m}) from the sequence $(b_{n_j, k_j})_j$ and setting here $\lambda(t) := \lim_{m \rightarrow \infty} b_{j_m}$, we obtain

$$\gamma(tx + (1-t)y) \leq \lambda(t)\gamma(x) + (1-\lambda(t))\gamma(y),$$

which was to be shown.

Note that in case (ii), with the same notations and the definition of the function λ , applying Lemma 5.1(ii), in a similar way, we first obtain the inequality

$$\gamma(a_{n,k}x + (1-a_{n,k})y) \geq b_{n,k}\gamma(x) + (1-b_{n,k})\gamma(y)$$

for all $x, y \in I, x < y$, and $k, n \in \mathbb{N}, k \leq n$, and then

$$\gamma(tx + (1-t)y) \geq \lambda(t)\gamma(x) + (1-\lambda(t))\gamma(y)$$

for $t \in (0, 1)$.

Part (iii) follows from parts (i) and (ii). □

In the next basic result we use parts (i) and (iii) of Lemma 5.4.

Lemma 5.5. *Let $I \subset \mathbb{R}$ be an interval. If a function $\gamma \in \mathcal{CM}_+(I)$ is such that for every $t \in (0, 1)$ there exists $\lambda(t) \in (0, 1)$ such that for all $x, y \in I, x \leq y$,*

$$\gamma(tx + (1-t)y) \leq \lambda(t)\gamma(x) + (1-\lambda(t))\gamma(y);$$

and for all $x, y \in I, x \geq y$,

$$\gamma(tx + (1-t)y) \geq \lambda(t)\gamma(x) + (1-\lambda(t))\gamma(y),$$

then there are $a, b \in \mathbb{R}, a \neq 0$, such that

$$\gamma(x) = ax + b, \quad x \in I.$$

Proof. Take $t = \frac{1}{2}$ and arbitrary $x, y \in I, x < y$. In view of the first assumed inequality we have

$$\gamma\left(\frac{x+y}{2}\right) \leq \lambda\left(\frac{1}{2}\right)\gamma(x) + \left(1-\lambda\left(\frac{1}{2}\right)\right)\gamma(y), \quad (11)$$

and from the second inequality (interchanging the roles of x and y)

$$\gamma\left(\frac{y+x}{2}\right) \geq \lambda\left(\frac{1}{2}\right)\gamma(y) + \left(1-\lambda\left(\frac{1}{2}\right)\right)\gamma(x). \quad (12)$$

It follows that

$$\lambda\left(\frac{1}{2}\right)\gamma(y) + \left(1-\lambda\left(\frac{1}{2}\right)\right)\gamma(x) \leq \lambda\left(\frac{1}{2}\right)\gamma(x) + \left(1-\lambda\left(\frac{1}{2}\right)\right)\gamma(y),$$

whence

$$\left(2\lambda\left(\frac{1}{2}\right) - 1\right)(\gamma(y) - \gamma(x)) \leq 0.$$

Since γ is increasing, we conclude that

$$\lambda\left(\frac{1}{2}\right) \leq \frac{1}{2}.$$

Similarly, for arbitrary $x, y \in I, x > y$, from the second assumed inequality we have

$$\gamma\left(\frac{x+y}{2}\right) \geq \lambda\left(\frac{1}{2}\right) \gamma(x) + \left(1 - \lambda\left(\frac{1}{2}\right)\right) \gamma(y), \quad (13)$$

and from the first inequality (where x and y are interchanged) we get

$$2\gamma\left(\frac{y+x}{2}\right) \leq \lambda\left(\frac{1}{2}\right) \gamma(y) + \left(1 - \lambda\left(\frac{1}{2}\right)\right) \gamma(x). \quad (14)$$

whence

$$\lambda\left(\frac{1}{2}\right) \gamma(x) + \left(1 - \lambda\left(\frac{1}{2}\right)\right) \gamma(y) \leq \lambda\left(\frac{1}{2}\right) \gamma(y) + \left(1 - \lambda\left(\frac{1}{2}\right)\right) \gamma(x),$$

which can be written in the form

$$\left(2\lambda\left(\frac{1}{2}\right) - 1\right) (\gamma(x) - \gamma(y)) \leq 0.$$

Since γ is strictly increasing we obtain that

$$\lambda\left(\frac{1}{2}\right) \geq \frac{1}{2},$$

and consequently

$$\lambda\left(\frac{1}{2}\right) = \frac{1}{2}.$$

Now inequalities (11), (12), (13) and (14) imply that

$$\gamma\left(\frac{x+y}{2}\right) = \frac{\gamma(x) + \gamma(y)}{2}, \quad x, y \in I,$$

that is γ is Jensen affine. The strict monotonicity of γ implies that $\gamma(x) = ax + b (x \in I)$, for some $a, b \in \mathbb{R}, a \neq 0$. This completes the proof. \square

Remark 5.6. Clearly, the counterparts of this lemma in all the remaining cases remain true.

6. Main results

Theorem 6.1. *Let $I \subset \mathbb{R}$ be an interval and $p, q, r, s \in [0, 1]$ such that $p + q \neq 1 \neq r + s$. Suppose that $\gamma \in \mathcal{CM}(I)$ and the functions $K, L : I^2 \rightarrow I$ are means in I .*

Then $L_{id}^{(r,s)} = K_{\gamma}^{(p,q)}$ if, and only if, $\gamma \sim id$ and

$$L(x, y) = \frac{(p-r)x + (q-s)y + (1-p-q)K(x, y)}{1-r-s}, \quad x, y \in I. \quad (15)$$

Proof. Without any loss of generality we can assume that γ is increasing. Assume that $L_{id}^{(r,s)} = K_{\gamma}^{(p,q)}$. Of course the numbers $p, q, r, s \in [0, 1]$ such that $p+q \neq 1 \neq r+s$ must satisfy the assumptions of exactly one of Lemmas 4.1–4.4. Applying this lemma together with Lemmas 5.4 and 5.5, we conclude that there are $a, b \in \mathbb{R}, a \neq 0$, such that

$$\gamma(x) = ax + b, \quad x \in I,$$

that is $\gamma \sim id$. Setting this function γ into equality $L_{id}^{(r,s)} = K_{\gamma}^{(p,q)}$ we obtain

$$rx + sy + (1-r-s)L(x, y) = px + qy + (1-p-q)K(x, y), \quad x, y \in I,$$

which implies (15). Since the converse implication is obvious, the proof is complete. \square

Hence, in view of Remark 3.2, we obtain

Theorem 6.2. *Let $I \subset \mathbb{R}$ be an interval, $K, L : I^2 \rightarrow I$ be means in $I, p, q, r, s \in [0, 1]$ be such that $p+q \neq 1 \neq r+s$, and $\varphi, \psi \in \mathcal{CM}(I)$. Then*

$$L_{\psi}^{(r,s)} = K_{\varphi}^{(p,q)}$$

if, and only if, $\psi \sim \varphi$ and

$$L(x, y) = \varphi^{-1} \left(\frac{(p-r)\varphi(x) + (q-s)\varphi(y) + (1-p-q)\varphi(K(x, y))}{1-r-s} \right)$$

for all $x, y \in I$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- [1] Daróczy, Z.: On the equality and comparison problem of a class of mean values. *Aequat. Math.* **81**, 201–208 (2011)
- [2] Daróczy, Z., Dašcal, J.: On the equality problem of conjugate means. *Results Math.* **58**, 69–79 (2010)
- [3] Daróczy, Z., Páles, Zs.: On means that are both quasi-arithmetic and conjugate arithmetic. *Acta Math. Hungar.* **90**, 271–282 (2001)
- [4] Daróczy, Z., Páles, Zs.: Gauss-composition of means and the solution of the Matkowski-Sutó problem. *Publ. Math. Debrecen* **61**(1–2), 157–218 (2002)
- [5] Maksa, Gy., Páles, Zs.: Remarks on the comparison of weighted quasi-arithmetic means. *Colloquium Math.* **120**, 77–84 (2010)
- [6] Matkowski, J.: Invariant and complementary means. *Aequat. Math.* **57**, 87–107 (1999)

Janusz Matkowski

Faculty of Mathematics, Computer Science and Econometrics

University of Zielona Góra

Szafrana 4a, 65-516 Zielona Góra, Poland

e-mail: J.Matkowski@wmie.uz.zgora.pl

Received: July 6, 2012

Revised: October 1, 2012